Brain-body dance: addressing mental health during injury rehabilitation

William Bracewell

My career in ballet is probably one that echoes many others. I started dancing around the age of 8, and was recognised by my teacher at the time as having potential to progress further. At the age of 11, I moved from Swansea to London, after receiving a place to live and train full time at The Royal Ballet School. This was an intense introduction into the ballet world to say the least; a pressure cooker environment that suited some and not others. At the time, there was a focus on hard training of the ballet skill, with less concern over the holistic development of the student. After 8 years of training I joined Birmingham Royal Ballet and danced professionally with the company for 7 years, touring nationally and internationally for most months of the year (sometimes for several months at a time). In 2017, I moved to The Royal Ballet, where I now dance as a Principal of the company. This means I perform the main or title roles in up to 13 productions each season.

SPINAL SURGERY

My injury history had been sprinkled with minor ankle issues and growth-related problems, but in April 2019 I underwent microdiscectomy surgery on my L5-S1 level in my spine. After a herniated disc injury 3 years prior, I had been managing ongoing back pain, and after what I initially thought was a minor hamstring tear, it quickly became apparent it was much more serious and now restricting the sciatic nerve in my left leg. I could not move at all without intense searing pain which immobilised me for weeks. I began a very slow and conservative rehabilitation approach, but 6 weeks into the programme, my leg would intermittently give way during the most basic warm-up dance movements, and I would end up collapsed on the floor.

At this point, I discussed my options with our clinical director: continue a conservative rehabilitation with the potential for surgery a year later, or opt for surgery immediately. The MRI suggested I had quite a severe herniation, which, coupled with the fact that I had been struggling with the rehabilitation

mentally—including ending up in tears on the floor during coaching sessions—made the surgical route seem like a clear choice. The healthcare team supported this decision, and while there was, of course, a large amount of trepidation ahead of spinal surgery, I'm happy that 5 years later it seems to have been the right decision.

CONNECTING MIND AND BODY

I would describe myself as incredibly ambitious, so to have extended periods of stagnant time after my surgery was very frustrating. I tried to fill my time with other interests that were not physical, but I missed the rigour and passion of dancing. Being back in the gym, sweating and working towards my goal of returning to the stage was a huge relief mentally. My physical rehab went well. I tried to crosspollinate the expertise of my rehab team

(figure 1) as much as possible; my Pilates and strength and conditioning (S&C) coaches would watch ballet coaching sessions to assess posture and consider new exercises that would be beneficial. Incorporating exercises that I could clearly associate with dance helped transfer the strength I gained during S&C and Pilates into the studio, which I otherwise had a tendency to disassociate. In many ways rehab suited my personality really well: I could schedule my days as I wanted; I could rest when I needed; and there were clear goals to be achieved. I cannot overemphasise how aware I became of the link between my mental and physical health. It seems obvious, but it's easy to detach the

ADDRESSING MY MENTAL HEALTH

An aspect I had not considered until it was looking me directly in the face was the potential loss of my profession and a huge majority of my life. I love my work, and the thought of that being taken away before I was ready was a difficult thing to process. There were also points early in the rehabilitation where I began to fear the



Figure 1 Celebrating with my rehab team following a successful return to the stage (from left to right: Gemma Hilton, Brian Maloney, William Bracewell, Jane Paris, Adam Mattiussi).

The Royal Ballet, Royal Opera House, London, UK



Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

Patient voices

loss of more general physical functions. For example, walking a dog that might pull in erratic ways, or bending to put on a sock comfortably. These fears eventually faded as my movement improved, but the initial worry was vivid. During this time I saw our in-house performance psychologist, who helped me work through the day-to-day mental difficulties, and identified elements of post-traumatic stress. With the advice of our wonderful internal healthcare department-and funding from a generous individual-I saw a psychotherapist. This became a turning point as we discussed life from a wider lens and assessed why I was putting myself through such an extreme process to continue dancing. I realised there was no other option than to give everything I could to try to return to dance. I would not be able to walk away from the situation without knowing I'd done everything in my power to get back to the stage. We assessed why I love dancing so much and why it was worth the effort. I learned that it's the effort that I love: I love to work, which in some ways was the reason I had become injured initially... by overexerting myself without adequate recovery.

MOVING MINDFULLY FORWARD

I wanted to highlight the mental health aspect of rehabilitation as it's maybe not

always factored into a programme but in my opinion was a key pillar in getting me back to full capacity. I have come to realise that strength and mobility work will be a part of my life forever, and I relish that. My fundamentals will be keeping a strong and mobile core and hips and moving as much as possible for as long as possible. I am aware that the neurological connection and sensitivity to that area of my back remains heightened 5 years later. I have had two subsequent MRI scans on my back after being worried something had happened again due to pain. This will probably be something I do every few years while I'm dancing to keep my mind at ease as much as keeping an eve on the condition of the disc. I still worry that I might injure my back again, not so much that it stops me dancing but there are some movements that still scare me and probably always will.

With high-profile athletes becoming more vocal about their mental health capacity and needs, I feel positive that this will become commonplace in dance rehabilitation.

I would encourage practitioners to value mental health as highly as strength training, mobility training and other physical therapies:

- ► Anticipate psychological challenges and programme therapy into a rehab plan.
- ► Promote honest discussion around mental health and function.
- Explore cross-pollination of coaches' expertise through joint sessions.

X William Bracewell @will_bracewell

Acknowledgements I would like to thank Adam Mattiussi, Jane Paris, Gemma Hilton, Brian Maloney, Greg Retter and Angela Bernstein for their support in helping me back to the stage.

Collaborators Joseph Shaw: The Royal Ballet, Royal Opera House, UK

Contributors Content of submission: WB. Support with editing and submission: JS.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; internally peer reviewed.

© Author(s) (or their employer(s)) 2024. No commercial re-use. See rights and permissions. Published by BMJ.



To cite Bracewell W. *Br J Sports Med* 2024;**58**:747–748.

Accepted 18 April 2024 Published Online First 7 June 2024

Br J Sports Med 2024;**58**:747–748. doi:10.1136/bjsports-2024-108471