# Paediatric post-concussive symptoms: symptom clusters and clinical phenotypes

Todd W Lyons <sup>(D)</sup>, <sup>1,2</sup> Rebekah Mannix, <sup>1,2</sup> Ken Tang, <sup>3</sup> Keith Owen Yeates, <sup>4,5</sup> Gurinder Sangha, <sup>6,7</sup> Emma CM Burns, <sup>8,9</sup> Darcy Beer, <sup>10</sup> Alexander S Dubrovsky, <sup>11,12</sup> Isabelle Gagnon, <sup>13</sup> Jocelyn Gravel, <sup>14</sup> Stephen B Freedman, <sup>15</sup> William Craig, <sup>16</sup> Kathy Boutis, <sup>17</sup> Martin H Osmond, <sup>3,18</sup> Gerard Gioia, <sup>19</sup> Roger Zemek <sup>(D)</sup>, <sup>3,18</sup> The Pediatric Emergency Research Canada (PERC) 5P Concussion Team<sup>20</sup>

## ABSTRACT

**Objective** To assess the co-occurrence and clustering of post-concussive symptoms in children, and to identify distinct patient phenotypes based on symptom type and severity.

**Methods** We performed a secondary analysis of the prospective, multicentre Predicting and Preventing Post-concussive Problems in Pediatrics (5P) cohort study, evaluating children 5–17 years of age presenting within 48 hours of an acute concussion. Our primary outcome was the simultaneous occurrence of two or more persistent post-concussive symptoms on the Post-Concussion Symptom Inventory at 28 days post-injury. Analyses of symptom and patient clusters were performed using hierarchical cluster analyses of symptom severity ratings.

**Results** 3063 patients from the parent 5P study were included. Median age was 12.1 years (IQR: 9.2-14.6 years), and 1857 (60.6%) were male. Fatigue was the most common persistent symptom (21.7%), with headache the most commonly reported co-occurring symptom among patients with fatigue (55%; 363/662). Headache was common in children reporting any of the 12 other symptoms (range: 54%-72%). Physical symptoms occurred in two distinct clusters: vestibularocular and headache. Emotional and cognitive symptoms occurred together more frequently and with higher severity than physical symptoms. Fatigue was more strongly associated with cognitive and emotional symptoms than physical symptoms. We identified five patient groups (resolved/minimal, mild, moderate, severe and profound) based on symptom type and severity. **Conclusion** Post-concussive symptoms in children occur in distinct clusters, facilitating the identification of distinct patient phenotypes based on symptom type and severity. Care of children post-concussion must be comprehensive, with systems designed to identify and treat distinct post-concussion phenotypes.

► Additional supplemental material is published online only. To view, please visit the journal online (http://dx.doi. org/10.1136/bjsports-2021-105193).

For numbered affiliations see end of article.

## Correspondence to

Dr Todd W Lyons, Departments of Pediatrics and Emergency Medicine, Harvard Medical School, Boston, USA; Todd.Lyons@childrens.harvard. edu

Accepted 18 February 2022 Published Online First 10 March 2022

## Check for updates

## © Author(s) (or their employer(s)) 2022. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Lyons TW, Mannix R, Tang K, *et al. Br J Sports Med* 2022;**56**:785–791.

**BMJ** 

## INTRODUCTION

Paediatric concussions are a major public health concern.<sup>1–5</sup> While the majority of children with a concussion will recovery completely within 4 weeks, nearly 30% will experience persistent post-concussive symptoms lasting 1 month or longer.<sup>6–11</sup> Prolonged symptoms can impact academic performance, participation in extra-curricular activities, and are associated with a lower quality of life.<sup>12–14</sup> Post-concussive symptoms are commonly

categorised into physical, cognitive, emotional and sleep domains.<sup>15 16</sup> However, little is known about the relationships between individual symptoms, the co-occurrence of symptoms in each of these domains, or whether specific clinical phenotypes can be identified based on symptom clusters. By understanding the relationships between post-concussive symptoms and identifying distinct clinical phenotypes in children with concussion, clinicians will be better prepared to identify and address the entirety of patients' post-concussion symptomatology, which may reduce time to recovery.

We sought to evaluate the nature of persistent post-concussive symptoms in a prospective cohort of children diagnosed with concussion in the emergency department (ED). We sought to define the associations between symptoms persisting 28 days following concussion and between symptom domains (physical, cognitive and emotional). Finally, we sought to identify novel patient phenotypes based on symptom type and severity.

## METHODS Study population

This was a secondary analysis of the Predicting and Preventing Post-concussive Problems in Pediatrics (5P) study.<sup>8</sup><sup>17</sup> We evaluated patients from both the derivation and validation cohorts. Patients were included if they were 5-17 years old, had a concussion defined by the Zurich consensus statement,<sup>10</sup> suffered their head injury within 48-hours of ED presentation and were proficient in English or French.<sup>17</sup> Patients were excluded if they had a Glasgow Coma Score of ≤13, abnormal neuroimaging, required neurosurgery, intubation or intensive care, or experienced multisystem injuries opmental delay with communication difficulties, children with no clear history of dren with a r dren with a previous history of enrolment in the study were also excluded.<sup>17</sup> Patients were screened and approached by research staff, and if willing were consented for study involvement.

## Study design/setting

The parent study was a prospective cohort study from August 2013 to June 2015 at 9 Canadian



## **Original research**

paediatric hospitals in the Pediatric Emergency Research Canada Network.<sup>17</sup> All centres are academic, paediatric centres with a combined annual volume of approximately 500000 ED visits. Neither patient nor public were involved in the design, conduct, reporting or dissemination of our research.

## **Data collection**

After obtaining informed consent and assent, as appropriate, parents completed the Acute Concussion Evaluation (ACE), a validated tool to objectively diagnose concussion (as indicated by  $\geq$ 1 symptoms on the ACE).<sup>18</sup> The following data were obtained from parents: demographics, medical history, presenting history and physical examination findings. Participant-reported ratings of current and pre-injury symptoms were obtained using the validated and reliable Post-Concussion Symptom Inventory (PCSI), where patients reported the presence and severity of each symptom relative to their pre-injury baseline.<sup>15</sup>

## Patient follow-up

Research assistants contacted participating families at 4 weeks after their index ED visit. Automated follow-up surveys using the Research Electronic Data Capture (REDCap)<sup>19</sup> data collection tool and telephone follow-up survey were used to collect patient self-reported symptoms via the PCSI.

## **Outcome measures**

Our primary outcome was the simultaneous occurrence of two or more post-concussive symptoms on the PCSI at 28 days post-injury. We chose to evaluate the 13 PCSI items common across all age groups (online supplemental table 1). PCSI scores for children aged 5-12 years, reported on a 0-2 scale, were multiplied by 3, so all scores were reported on the same (0-6) ordinal scale.<sup>20</sup> For initial analyses examining the co-occurrence of symptom pairs, we defined a post-concussive symptom as a dichotomous outcome (any positive difference between a patient-reported symptom at 28 days post-concussion minus their pre-injury symptom rating). For hierarchical cluster analyses, we analysed both the presence and severity (magnitude of the delta between pre-injury symptom rating score and patientreported symptom severity) of symptoms.

## **Statistical analyses**

We described the patient population using medians and interquartile ranges (IQRs) for continuous variables, and frequency and percentages for categorical variables. We analysed the relationship between post-concussion symptoms, describing their co-occurrence using conditional percentages (the portion of patients with symptom X who also have symptom Y). All usable PCSI data, including those from partially completed questionnaires, were included in this analysis.

We next analysed the relationship between post-concussive symptoms using hierarchal cluster analysis, reflecting clustering of postconcussive symptoms by type and severity. Because not all patients had data for all PCSI outcomes, we excluded those missing any PCSI data. We compared characteristics of included and excluded patients to ensure there were no significant differences.

To identify symptom and patient clusters (clinical phenotypes), hierarchical agglomerative cluster analysis (HCA) was applied in both dimensions to evaluate interrelationships among the 13 PCSI items, and to examine variations among patients based on their symptom profiles (ie, response patterns across PCSI items). Because of a relatively large number of unique patient symptom profiles, prior to HCA, an initial k-means clustering was performed on the patient

dimension to reduce the number of observations to a manageable size (40 patient clusters).<sup>21</sup> Each of the 40 clusters contained varying numbers of patients that share similar symptom profiles. Cluster means were then computed (for each PCSI item) and applied to our HCAs using Ward's dissimilarity-based agglomerative algorithm (minimum variance method), where clustering criterion is based on squared Euclidean distances.<sup>22</sup>

To present our findings, two sets of dendrograms (tree diagrams) accompanied by an associated data heatmap were constructed. The two dendrograms represent an empirically derived 'classification' of symptoms and patients, respectively, Protected based on the extent of (dis)similarity among constituents. The resulting heatmap provides an overview of study data summarised at the level of the initial patient clusters (n=40). 'Cutting' the dendrogram was performed to facilitate symptom and patients by copyright groups at a level of precision, which was determined a priori. For symptoms groups (composed of the 13 analysed symptoms), the dendrogram was 'cut' (based on tree height) to facilitate the emergence of four distinct groups, which was chosen to match the number of theoretical domains (physical, emotional, cogni-; including tive and fatigue) that originally conceptualised the PCSI.<sup>15</sup> Likewise, because of the large number of unique patient symptom profiles (40), prior to HCA, an initial k-means clustering was performed on patient dimension. We elected to 'cut' the patient dendrogram to identify five distinct patient groups. uses related to text

The study's sample size was determined by the number of patients in the parent 5P study. All analyses were performed using R version  $4.0.3.^{24}$ 

## RESULTS

## **Patient population**

Screening was performed on 8046 patients (5229 from the deriand data mining vation cohort and 2817 from the validation cohort), of whom 3063 (38%) were enrolled; all of whom we included. Median patient age was 12.1 years (IOR: 9.2-14.6 years), and 1857 (60.6%) were male. Sports and recreational injuries accounted for the majority (68.1%) of included concussions, followed by falls (28.6%). Characteristics of the included population are summarised in table 1. For purposes of evaluating symptom I, AI training, and similar and patient clusters, 2355 (76.9%) patients without any missing data from the PCSI at 28 days post-concussion were included (table 1). Differences between included and excluded patients in this analysis are summarised in online supplemental table 2.

## **Co-occurrence of post-concussive symptoms**

The co-occurrence of post-concussive symptoms 28 days post-injury is described in table 2. Fatigue was the most commonly reported symptom (21.7%) and was reported as a concurrent symptom in 62%–76% of patients reporting any other post-concussive symptom. Among patients reporting fatigue (N=665), only headache was reported in more than 50% of patients. Headache was the second most commonly reported symptom (19.3%) and was reported as a concurrent symptom in 54%-72% of patients reporting any other post-concussive symptom. Among patients with headache (N=593), fatigue (62%) and difficulty concentrating (50%) were the most commonly reported co-occurring post-concussive symptoms. Difficulty concentrating was reported by 488 (15.9%) patients and was reported as a concurrent symptom in 49%–72% of patients reporting any other post-concussive symptom. Vision problems were the least commonly reported post-concussive symptom (6.7%) but occurred with a high frequency (39%) among those with balance problems, and were present among 22%-35% of patients reporting any other post-concussive symptom.

technologies

ę

Table 1     Patient chara	acteristics of included child	Iren with a concussion
Characteristic	Patients included in bivariate symptom analyses N=3063	Patients included in hierarchical symptom analyses N=2553
Age (years), median (IQR)	12.0 (9.2–14.6)	12.0 (9.3–14.6)
Male sex, n/total (%)	1857/3062 (60.6%)	1416/2354 (60.2%)
Time from Injury to ED triage (hours), median (IQR)	2.9 (1.5–11.3)	3.0 (1.5–12.6)
Mechanism of injury, n/ total (%)		
Sports/recreation	2075/3049 (68.1%)	1614/2353 (68.6%)
Fall	871 (28.6%)	659 (28.0%)
Motor vehicle crash	57 (1.9%)	45 (1.9%)
Assault	43 (1.4%)	32 (1.4%)
Other	3 (0.1%)	3 (0.1%)
Loss of consciousness, n/ total (%)	395/2712 (14.6%)	304/2084 (14.6%)
Post-injury seizure, n/ total (%)	57/3041 (1.9%)	41/2348 (1.7%)
Glasgow Coma Score, median (IQR)	15 (15–15)	15 (15–15)
History, n/total (%)		
Prior treatment for headache	518/3049 (17.0%)	400/2353 (17.0%)
Personal history of migraine	392/3038 (12.9)	310/2342 (13.2%)
Learning disability	243/3039 (8.0%)	179/2347 (7.6%)
Attention deficit hyperactivity disorder	268/3036 (8.8%)	195/2345 (8.3%)
Anxiety	237/3045 (7.8%)	188/2349 (8.0%)
Depression	87/3047 (2.9%)	62/2350 (2.6%)
Sleep disorder	62/3040 (2.0%)	47/2348 (2.0%)
Other psychiatric disorders	32/3016 (1.1%)	24/2327 (1.0%)
Family history of migraine, n/total (%)	1436/2981 (48.2%)	310/2342 (13.2%)
BESS number of tandem stance errors, median (IQR)	3 (1–8)	3 (1–8)
BESS number of double leg stand errors, median (IQR)	0 (0–1)	0 (0–1)
PCSI scores at index ED visit, median (IQR)		
Physical	1.5 (0.9–2.5)	1.5 (0.9–2.4)
Fatigue	2.0 (0.7–3.7)	2.0 (0.7–3.7)
Emotional	0.5 (0.0–1.5)	0.5 (0.0–1.5)
Cognitive	0.8 (0.0–2.0)	0.6 (0.0–2.0)
PCSI scores represent item a	averages for each domain.	

BESS, Balance Error Scoring System; ED, emergency department; PCSI, Post-

Concussion Symptom Inventory.

## Clusters and phenotypes of post-concussive symptoms

Analysis of patient and symptom clusters by symptom severity using HCA is seen in figure 1. Overall, this analysis revealed that physical, emotional, cognitive and fatigue symptoms clustered together, consistent with the proposed structure of the PCSI. Physical symptoms further differentiated into a vestibularocular cluster (balance problems, dizziness, vision problems and nausea) and a headache cluster (headache, sensitivity to light and sensitivity to noise). Fatigue was more closely related to cognitive and emotional symptoms than to physical symptoms.

The k-means analysis identified the initial 40 clusters of post-concussive patients. The largest cluster of patients (1371 (44.8%)) was asymptomatic at 28 days. The next largest cluster (96 (4.1%)) reported mild fatigue as their only post-concussive symptom. One cluster of 9 (0.4%) children reported a high symptom burden across all domains. Overall, the analysis yielded five broad groups of clusters that varied by symptom type and severity (table 3). Group 1 (clusters 1-17) was fully recovered or were minimally symptomatic. Group 2 (clusters 18-23) was mildly symptomatic, but had higher symptom burden in the cognitive/emotional and fatigue domains. Group 3 (clusters 24–32) was moderately symptomatic, and was further divided into those with a higher burden of balance, dizziness and head-ache symptoms (clusters 24–27) and those with more headache, ŝ cognitive and fatigue symptoms (clusters 28-32). Group 4 (clusters 33-35) was highly symptomatic, largely across cognitive, copyright emotional and fatigue domains, with minimal physical symptoms. Group 5 (clusters 36-40) was profoundly symptomatic with high symptom burden across all domains, with only one cluster of patients (cluster 36) reporting low symptom burden in the vestibular-ocular domain. Clusters of patients reporting a high burden of vestibular-ocular symptoms (clusters 38–40) had the highest cumulative symptom burden. Overall, symptom severity was higher in clusters of patients reporting cognitive and emotional symptoms than in those reporting physical symptoms.

## DISCUSSION

In this multicentre, prospective study of children with concussion, we found important overlap between post-concussive symptoms at 28 days and describe novel clinical phenotypes of post-concussion patients. Fatigue was the most common persistent post-concussive symptom and was more closely related to cognitive and emotional symptoms than to physical symptoms, with implications for how this common symptom is treated. Additionally, we found that physical symptoms divided into vestibular-ocular and headaches clusters, while cognitive and emotional symptoms clustered together, and were associated with a higher symptom burden. Finally, we identified novel patient phenotypes and groups of post-concussive patients based on symptom type and severity. Together, these results support the need for clinicians to assess the entirety of a patient's symptomatology, and identify phenotypes of patients who may require distinct treatments. Furthermore, these data underscore the importance of multidisciplinary treatment approaches, with the goal of accelerating time to recovery.

While our post-concussion patient phenotypes were identified quantitatively, they reflect previously reported patient groupings.<sup>25-33</sup> A prior analysis evaluating symptoms in adolescents identified four novel symptom clusters, including cognitivefatigue-migraine, affective, somatic and sleep.<sup>26</sup> Another analvsis in a sports medicine clinic identified 3 clusters of symptoms: neurocognitive, somatic and emotional, with a higher burden of emotional symptoms.<sup>28</sup> Finally, the headache, vestibular-ocular and emotional symptom clusters we identified are congruent with other proposed symptom classification phenotypes.<sup>25 29 34 35</sup> Our study augments these previous works through its large sample size, prospective design, and incorporation of both the presence and severity of symptoms. Finally, this work builds on our prior latent class analyses of the 5P data, which identified distinct phenotypes of acute parent-reported symptoms, and assessed the association between these phenotypes and the diagnosis of postconcussion symptoms.<sup>36</sup> It is still not known how these acute

ð

uses related

to text

and

data mining

, AI training,

, and

simila

technologies.

# Br J Sports Med: first published as 10.1136/bjsports-2021-105193 on 10 March 2022. Downloaded from http://bjsm.bmj.com/ on May 31, 2025 at Department GEZ-LTA Erasmushogeschool . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

# **Original research**

Table 2 Bivaria	te comparisor	n of the prese	ence of indivi	idual post-con	icussion sympto	oms at 28 days							
Symptoms, N (%) of total patients reporting this symptom	Headache	Nausea	Problem with balance	Dizziness	Sensitivity to light	Sensitivity to noise	Vision problems	Fatigue	Irritability	Sadness	Nervousness	Difficulty concentrating	Difficulty remembering
Headache, N=593 (19.4%)	100%	34%	33%	44%	45%	46%	25%	62%	41%	27%	28%	50%	37%
Nausea, N=296 (9.7%)	68%	100%	43%	53%	48%	51%	31%	68%	52%	35%	36%	53%	42%
Problem with balance, N=294 (9.6%)	67%	43%	100%	68%	55%	56%	39%	71%	53%	41%	37%	65%	42%
Dizziness, N=358 (11.7%)	72%	44%	56%	100%	52%	54%	35%	72%	50%	38%	34%	62%	46%
Sensitivity to light, N=382 (12.5%)	%69	37%	43%	49%	100%	64%	32%	68%	49%	34%	34%	61%	45%
Sensitivity to noise, N=412 (13.5%)	66%	37%	40%	47%	60%	100%	30%	68%	53%	32%	37%	63%	47%
Vision problems, N=205 60.7%)	70%	45%	56%	60%	60%	60%	100%	75%	50%	43%	42%	68%	55%
Fatigue, N=665 (21.7%)	55%	30%	32%	39%	39%	42%	23%	100%	47%	31%	30%	49%	36%
Irritability, N=462 (15.1%)	54%	33%	34%	39%	41%	47%	22%	68%	100%	41%	38%	55%	41%
Sadness, N=272 (8.9%)	60%	38%	44%	50%	48%	47%	32%	76%	%69	100%	53%	60%	48%
Nervousness, N=276 (8.9%)	59%	39%	40%	45%	47%	55%	31%	73%	64%	52%	100%	62%	47%
Difficulty concentrating, N=488 (15.9%)	61%	32%	38%	45%	48%	53%	29%	67%	52%	33%	35%	100%	52%
Difficulty remembering, N=354 (11.6%)	62%	35%	45%	47%	48%	53%	32%	68%	52%	37%	36%	72%	100%
Percentages represen Colours correspond to	of the proportion	of patients wit of co-occurence	th the symptom of post-concus	n in column 1 wh. ssion symptoms.	o are concurrently ranging from gree	reporting the add	itional symptoms	in the top rov	v. ro-occurence and	I red the highes	tt co-occurence of s	umntoms	



Figure 1 Clusters of post-concussive symptoms in paediatric patients.

patient symptom phenotypes predict patient symptomatology and phenotypes at 28 days post-concussion.

# Symptom co-occurrence, clusters and novel patient phenotypes

We uncovered important findings regarding the co-occurrence of post-concussive symptoms. Fatigue and headache were the most commonly described symptoms. However, among patients reporting fatigue or headache, few other symptoms were commonly reported. Furthermore, while vision problems were the least commonly reported symptom, those with vision problems had an overall higher symptom burden. These

Table 3	Groupings of patient phenotypes by symptom type and
severity	

Group	Clusters	Name	Description
1	1–17	Resolved or minimally symptomatic	No symptoms (cluster 1) or minimal symptoms (clusters 2–16) reported across all domains
2	18–23	Mildly symptomatic	Mildly symptomatic with higher symptoms severity in fatigue and cognitive emotional domains
3	24–32	Moderately symptomatic	Moderately symptomatic and further divided into: balance, dizziness and headache subgroup (clusters 24–27) and headache, cognitive and fatigue subgroup (clusters 28–32)
4	33–35	Highly symptomatic	Highly symptomatic but largely across cognitive/emotional and fatigue domains with minimal physical symptoms
5	36–40	Profoundly symptomatic	Profoundly symptomatic with a high symptom burden across all domains with exception of cluster 36 reporting few vestibular-ocular symptoms

data suggest some symptoms may be more likely to occur in isolation, while others, such as vision problems, sadness and problems with balance, may be an indicator of higher overall symptom burden. By being aware of symptoms which represent an overall high symptom burden, clinicians may identify those children more likely to benefit from earlier or more aggressive therapies or interventions. Furthermore, our results emphasise that while some symptoms may occur with less frequency than others, the lowest co-occurrence of symptoms (irritability and vision problems) still occurred together in more than 1 in 5 children, emphasising the importance of comprehensive symptom assessment.<sup>37</sup>

The clusters of patients and symptoms we identified have important implications for treating children following a concussion. First, and reassuringly, the largest group (group 1) of patients based on symptom type and severity was of those who were asymptomatic or minimally symptomatic at 28 days. Second, these results highlight the importance of addressing fatigue, the most common post-concussive symptom, which commonly coexists with other symptoms, and if not properly addressed could potentially prolong overall symptoms duration. Aetiologies for fatigue following concussion are multifactorial.<sup>38–40</sup> Addressing sleep issues is one possible focus for clinicians treating fatigue in concussed children.<sup>41–43</sup> Importantly, fatigue was more strongly associated with cognitive and emotional than physical symptoms, suggesting it may represent a mental rather than physical phenomenon by 4 weeks post-injury, with critical implications for how it is best treated.<sup>44 45</sup> Third, while moderately symptomatic patients were more troubled by headache, among those patients with severe symptom burden, the severity and burden of cognitive and emotional symptoms often exceeded those of physical symptoms. These results highlight the importance of recognising and treating the often difficult to manage cognitive and emotional post-concussion symptoms.<sup>46</sup> Fourth, we found that

## **Original research**

physical symptoms further differentiated into vestibular-ocular and headache clusters. While the most symptomatic patients may have higher severity symptoms in both physical domains, patients with more mild and moderate symptoms tend to have a predominance of either headache or vestibular-ocular symptoms but not both. Fifth, a high burden of vestibular-ocular symptoms was associated with an overall high symptom burden across all domains.<sup>47</sup> Previous literature has suggested therapies targeted at improving vestibular-ocular symptoms may improve patient outcomes.<sup>48</sup> Finally, among the most severely symptomatic children, the severity of vestibular-ocular, headache, cognitive, emotional and fatigue symptoms varied. These results demonstrate the heterogeneity in symptoms that are characteristic even among the most severely affected patients and underscore the need for clinicians to develop specific treatment strategies for individual patients.

## Implications for concussion care and research

This work has important implication for clinicians caring for children with post-concussion symptoms. These data inform clinicians about other symptoms to be assessed for when patients report a specific symptom. Additionally, by identifying novel phenotypes of patients, clinicians can understand how symptoms travel together. Failure to address co-occurring symptoms may lead to prolonged recovery, despite addressing the patient's primary complaint. Furthermore, the high burden of cognitive and emotional symptoms among the most severely symptomatic patients stresses the importance of a multidisciplinary, biopsychosocial approach to treatment.<sup>49</sup> In addition, future studies evaluating concussion and targeting interventions for specific post-concussion symptoms such as headache should take into account the co-occurrence of symptoms when designing outcome assessments.

## Methodologic considerations and limitations

Our study must be interpreted in the context of its limitations. First, all participants were recruited after presenting to the ED for evaluation, and may be more severely concussed than children presenting to other clinical settings. Therefore, our results may not be applicable to patients presenting with concussion to other clinical settings. However, recent data suggest this assumption about site of initial concussion care and sevrity of injury may not be correct.<sup>50 51</sup> Second, we were missing follow-up symptom data for some patients, requiring us to eliminate them from our HCA. However, we found few clinically meaningful differences between those patients for whom all data were available and those with missing data. Third, we included patients over a range of ages. Clinical phenotypes and co-occurrence of symptoms may vary by age and sex. Furthermore, we multiplied symptom scores for the youngest to normalise symptom scales. While this method has been previously reported, the validity of this approach is not known. Fourth, sleep problems are commonly reported post-concussion, but we could only report on fatigue, as it was the only common element across all age groups in PCSI. Fifth, we included any increase in symptom severity over baseline for our analysis of the co-occurrence. In doing so, a patient with even a minor increase from their pre-injury state would be classified similarly to a patient with a marked increase in symptomatology. However, this methodology has been previously described, 52-55 and the magnitude of symptom change on the PCSI was accounted for in our HCA. Finally, we only report patient clusters and phenotypes at 28 days post-injury. Prior data have suggested changes in the types and severity of symptoms

at various intervals post-recovery.<sup>11</sup> However, the early identification of clinical phenotypes could allow for targeted interventions. Further work is needed to understand how these symptom clusters and clinical phenotypes evolve over time as well as how patient-level factors such as age and gender impact symptom type and severity.

## **CONCLUSIONS**

In this prospective, multicentre study, we identified the frequency with which post-concussive symptoms occur together, Protected and described clusters of post-concussive symptoms and novel patient phenotypes at 28 days post-concussion. These data patient premotypes at 25 days post-concussion. These data underscore the need for comprehensive, multidisciplinary treatment programmes and individualised management plans based on symptomatology. They also highlight the need to address the high burden of cognitive, emotional and fatigue symptoms among the most symptomatic patients. Care systems designed to treat these distinct post-concussion phenotypes are needed, with the goal of reducing time to recovery.
Key messages
What is already known on this topic?
⇒ Post-concussive symptoms are commonly categorised into physical, cognitive, emotional and sleep domains.
What this study adds?
⇒ In this study of more than 3000 children with a concussion, the relationships between post-concussion symptoms at 28 days were examined, and novel phenotypes of patients with post-concussion symptoms were empirically defined.
How this study might affect research, practice or policy?
⇒ These data underscore the need for comprehensive, multidisciplinary treatment programmes and individualised management plans for patients following concussion.
Author affiliations
<sup>1</sup>Department of Pediatrics, Children's Hospital of Western Ontario, Lanada
<sup>2</sup>Department of Pediatrics, Children's Hospital of Western Ontario, Lanada
<sup>3</sup>Department of Pediatrics, Children's Hospital of Western Ontario, Lanada
<sup>3</sup>Department of Pediatrics, Children's Hospital of Western Ontario, Lanada
<sup>3</sup>Department of Pediatrics, Children's Hospital of Western Ontario, Lanada
<sup>3</sup>Department of Pediatrics, Children's Hospital of Western Ontario, Lanada
<sup>3</sup>Department of Pediatrics, Children's Hospital of Western Ontario, Lanada
<sup>3</sup>Department of Pediatrics, Children's Hospital Foundation of Manitoba, Winnipeg, Manitoba, Canada
<sup>3</sup>Departriment of Pediatrics, Montreal Cuebec, Canada
 underscore the need for comprehensive, multidisciplinary treatment programmes and individualised management plans based

<sup>12</sup>Department of Pediatrics, Montreal Children's Hospital - McGill University Health Centre, Montreal, Quebec, Canada

<sup>3</sup>Division of Physical and Occupational Therapy, McGill University Health Centre, Montreal, Quebec, Canada

<sup>1</sup>Department of Pediatrics, Saint Justine Hospital, Montreal, Quebec, Canada <sup>15</sup>Department of Pediatrics and Emergency Medicine, Cumming School of Medicine,

University of Calgary, Calgary, Alberta, Canada <sup>6</sup>Department of Pediatrics, Stollery Children's Hospital, Edmonton, Alberta, Canada

<sup>17</sup>Department of Pediatrics, The Hospital for Sick Children, Toronto, Ontario, Canada <sup>18</sup>Department of Pediatrics, University of Ottawa, Ottawa, Ontario, Canada

<sup>19</sup>Division of Neuropsychology, Children's National Hospital, Washington, District of Columbia USA

<sup>20</sup>Pediatric Emergency Research Canada, Calgary, Alberta, Canada

**Correction notice** This article has been corrected since it published Online First. The abstract has been amended.

**Acknowledgements** We are grateful to the parents and children who participated in this study. We appreciate the collaboration of the treating physicians and the research assistants at each participating site. We would like to thank the Pediatric Emergency Research Canada Network for making this study possible.

**Collaborators** For the Pediatric Emergency Research Canada 5P Concussion Team.

**Contributors** TWL, RM, KT and RZ designed the study and performed and interpreted data analyses. KY, GS, EB, DB, AD, WC, KB, MO, GG and RZ supervised data collection at the participating sites. TWL, RM and RZ drafted the manuscript. All authors critically reviewed and revised the manuscript and contributed to the final submission. TWL is responsible for the overall content as guarantor.

**Funding** This study was supported by operating grant (126197) and planning grant (MRP: #119829) from the Canadian Institutes of Health Research and grant (TM1:#127047) from the Canadian Institutes of Health Research–Ontario Neurotrauma Foundation Mild Traumatic Brain Injury Team. The funders played no role in the design and conduct of the study; collection, management, analysis or interpretation of the data; preparation, review or approval of the manuscript; and decision to submit the manuscript for publication.

**Competing interests** GG is an author of the Post-Concussion Symptom Inventory (PCSI) used in this study. The PCSI is freely available and he receives no financial benefit from its use.

Patient consent for publication Not applicable.

**Ethics approval** This study involves human participants. The ethics committees of each participating centre approved this study with permission for data sharing. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

**Data availability statement** Data are available upon reasonable request. Data for this manuscript are stored and available online in BrainCode and are available upon reasonable request.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

## ORCID iDs

Todd W Lyons http://orcid.org/0000-0002-7033-1070 Roger Zemek http://orcid.org/0000-0001-7807-2459

## REFERENCES

- Fridman L, Scolnik M, Macpherson A, et al. Annual trends in follow-up visits for pediatric concussion in emergency departments and physicians' offices. J Pediatr 2018;192:184–8.
- 2 National center for injury prevention. Report to Congress on mild traumatic brain injury in the United States: steps to prevent a serious public health problem. Atlanta, GA, 2003.
- 3 Veliz P, McCabe SE, Eckner JT, et al. Prevalence of concussion among US adolescents and correlated factors. JAMA 2017;318:1180–2.
- 4 Meehan WP, Mannix R. Pediatric concussions in United States emergency departments in the years 2002 to 2006. *J Pediatr* 2010;157:889–93.
- 5 Mannix R, O'Brien MJ, Meehan WP. The epidemiology of outpatient visits for minor head injury. *Neurosurgery* 2013;73:129–34.
- 6 Babcock L, Byczkowski T, Wade SL, et al. Predicting postconcussion syndrome after mild traumatic brain injury in children and adolescents who present to the emergency department. JAMA Pediatr 2013;167:156–61.
- 7 Barlow M, Schlabach D, Peiffer J, et al. Differences in change scores and the predictive validity of three commonly used measures following concussion in the middle school and high school aged population. Int J Sports Phys Ther 2011;6:150–7.
- 8 Zemek R, Barrowman N, Freedman SB, *et al*. Clinical risk score for persistent Postconcussion symptoms among children with acute concussion in the ED. *JAMA* 2016;315:1014–25.
- 9 Eisenberg MA, Andrea J, Meehan W, et al. Time interval between concussions and symptom duration. *Pediatrics* 2013;132:8–17.
- 10 NcCrory P, Meeuwisse W, Dvorak J. Consensus statement on concussion in sport—the 5 th International Conference on concussion in sport held in Berlin, October 2016. Br J Sports Med 2017;51.
- 11 Eisenberg MA, Meehan WP, Mannix R. Duration and course of Post-Concussive symptoms. *Pediatrics* 2014;133:999–1006.

- 12 Moran LM, Taylor HG, Rusin J, *et al*. Quality of life in pediatric mild traumatic brain injury and its relationship to postconcussive symptoms. *J Pediatr Psychol* 2012;37:736–44.
- 13 Novak Z, Aglipay M, Barrowman N, *et al*. Association of persistent Postconcussion symptoms with pediatric quality of life. *JAMA Pediatr* 2016;170:e162900.

**Original research** 

- 14 Yeates KO, Kaizar E, Rusin J, et al. Reliable change in postconcussive symptoms and its functional consequences among children with mild traumatic brain injury. Arch Pediatr Adolesc Med 2012;166:615–22.
- 15 Sady MD, Vaughan CG, Gioia GA. Psychometric characteristics of the postconcussion symptom inventory in children and adolescents. *Arch Clin Neuropsychol* 2014;29:348–63.
- 16 Gioia GA, Isquith PK, Schneider JC, et al. New approaches to assessment and monitoring of concussion in children. Top Lang Disord 2009;29:266–81.
- 17 Zemek R, Osmond MH, Barrowman N. Predicting and preventing postconcussive problems in paediatrics (5p) study: protocol for a prospective multicentre clinical prediction rule derivation study in children with concussion. *BMJ Open*2013;3:e003550.
- 18 Gioia GA, Collins M, Isquith PK. Improving identification and diagnosis of mild traumatic brain injury with evidence: psychometric support for the acute concussion evaluation. J Head Trauma Rehabil 2008;23:230–42.
- 19 Harris PA, Taylor R, Thielke R, et al. Research electronic data capture (REDCap)—A metadata-driven methodology and workflow process for providing translational research informatics support. J Biomed Inform 2009;42:377–81.
- 20 Teel EF, Zemek RL, Tang K, *et al*. The stability of retrospective Pre-injury symptom ratings following pediatric concussion. *Front Neurol* 2019;10.
- 21 Hartigan JA, Wong MA. Algorithm as 136: a k-means clustering algorithm. Appl Stat 1979;28:100.
- 22 Ward JH. Hierarchical grouping to optimize an objective function. *J Am Stat Assoc* 1963;58:236–44.
- 23 Murtagh F, Legendre P. Ward's Hierarchical Agglomerative Clustering Method: Which Algorithms Implement Ward's Criterion? *Journal of Classification* 2014;31:274–95.
- 24 R Core Team. R: a language and environment for statistical computing, 2020. Available: http://www.r-project.org/
- 25 Howell DR, Kriz P, Mannix RC, et al. Concussion symptom profiles among child, adolescent, and young adult athletes. Clin J Sport Med 2019;29:391–7.
- 26 Kontos AP, Elbin RJ, Schatz P, et al. A revised factor structure for the postconcussion symptom scale: baseline and postconcussion factors. Am J Sports Med 2012;40:2375–84.
- 27 Heyer GL, Young JA, Fischer AN. Lightheadedness after concussion: not all dizziness is vertigo. *Clin J Sport Med* 2018;28:272–7.
- 28 Joyce AS, Labella CR, Carl RL. The Postconcussion symptom scale: utility of a threefactor structure. *Med Sci Sports Exerc* 2015;47:1119–23.
- 29 Maruta J, Lumba-Brown A, Ghajar J. Concussion subtype identification with the Rivermead Post-concussion symptoms questionnaire. *Front Neurol* 2018;9:1034.
- 30 Howell DR, O'Brien MJ, Beasley MA, et al. Initial somatic symptoms are associated with prolonged symptom duration following concussion in adolescents. Acta Paediatr 2016;105:e426–32.
- 31 Guty E, Arnett P. Post-concussion symptom factors and neuropsychological outcomes in collegiate athletes. J Int Neuropsychol Soc 2018;24:684–92.
- 32 Langdon S, Königs M, Adang EAMC, *et al.* Subtypes of sport-related concussion: a systematic review and Meta-cluster analysis. *Sports Med* 2020;50:1829–42.
- 33 Lau BC, Collins MW, Lovell MR. Cutoff scores in neurocognitive testing and symptom clusters that predict protracted recovery from concussions in high school athletes. *Neurosurgery* 2012;70:371–9.
- 34 Lumba-Brown A, Ghajar J, Cornwell J, et al. Representation of concussion subtypes in common postconcussion symptom-rating scales. Concussion 2019;4:CNC65.
- 35 Craton N, Ali H, Lenoski S. Coach cv: the seven clinical phenotypes of concussion. *Brain Sci* 2017;7:119.
- 36 Yeates KO, Tang K, Barrowman N, et al. Derivation and initial validation of clinical phenotypes of children presenting with concussion acutely in the emergency department: latent class analysis of a multi-center, prospective cohort, observational study. J Neurotrauma 2019;36:1758–67.
- 37 Stillman A, Alexander M, Mannix R, et al. Concussion: evaluation and management. Cleve Clin J Med 2017;84:623–30.
- 38 Gagner C, Landry-Roy C, Lainé F, et al. Sleep-Wake disturbances and fatigue after pediatric traumatic brain injury: a systematic review of the literature. J Neurotrauma 2015;32:1539–52.
- 39 Riccardi JS, Ciccia A. Cognitive fatigue in pediatric traumatic brain injury: a metaanalysis and scoping review. J Head Trauma Rehabil 2021;36:226–41.
- 40 Crichton A, Anderson V, Oakley E, et al. Fatigue following traumatic brain injury in children and adolescents: a longitudinal follow-up 6 to 12 months after injury. J Head Trauma Rehabil 2018;33:200–9.
- Hinds A, Jungquist CR, Leddy JJ, et al. Sleep disturbance in patients with chronic concussive effects. Concussion 2016;1:CNC15.
- 42 Mosti C, Spiers MV, Kloss JD. A practical guide to evaluating sleep disturbance in concussion patients. *Neurology* 2016;6:129–37.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

# **Original research**

- 43 Kostyun RO, Milewski MD, Hafeez I. Sleep disturbance and neurocognitive function during the recovery from a sport-related concussion in adolescents. *Am J Sports Med* 2015;43:633–40.
- 44 Sharpe M. Abc of psychological medicine: fatigue. BMJ 2002;325:480-3.
- 45 Broshek DK, De Marco AP, Freeman JR. A review of post-concussion syndrome and psychological factors associated with concussion. *Brain Injury* 2015;29:228–37.
- 46 McCarty CA, Zatzick D, Stein E, et al. Collaborative care for adolescents with persistent Postconcussive symptoms: a randomized trial. *Pediatrics* 2016;138. doi:10.1542/peds.2016-0459. [Epub ahead of print: 13 09 2016].
- 47 Master CL, Master SR, Wiebe DJ, et al. Vision and vestibular system dysfunction predicts prolonged concussion recovery in children. *Clin J Sport Med* 2018;28:139–45.
- 48 Schneider KJ, Meeuwisse WH, Barlow KM, et al. Cervicovestibular rehabilitation following sport-related concussion. Br J Sports Med 2018;52:100–1.
- 49 Schneider KJ. Sport-Related concussion: optimizing treatment through evidenceinformed practice. J Orthop Sports Phys Ther 2016;46:613–6.

- 50 Zogg CK, Haring RS, Xu L, et al. The epidemiology of pediatric head injury treated outside of hospital emergency departments. *Epidemiology* 2018;29:269–79.
- 51 Risen SR, Reesman J, Yenokyan G, *et al.* The course of concussion recovery in children 6-12 years of age: experience from an interdisciplinary rehabilitation clinic. *PM&R* 2017;9:874–83.
- 52 Macnow T, Curran T, Tolliday C, *et al*. Effect of screen time on recovery from concussion. *JAMA Pediatr* 2021;175:1124.
- 53 Meehan WP, Mannix RC, Stracciolini A, et al. Symptom severity predicts prolonged recovery after sport-related concussion, but age and amnesia do not. J Pediatr 2013;163:721–5.
- 54 Meehan WP, O'Brien MJ, Geminiani E, et al. Initial symptom burden predicts duration of symptoms after concussion. J Sci Med Sport 2016;19:722–5.
- 55 Meehan WP, Mannix R, Monuteaux MC, et al. Early symptom burden predicts recovery after sport-related concussion. *Neurology* 2014;83:2204–10.

## Supplemental Table 1: PCSI Symptom Inventory By Age with Common Question 13 Common Items Across All Age Groups Highlighted

Age 5-7	Age 8-12	Age 13-18
	Common Elements Included	
Have you had headaches? Has your head hurt?	Have you had headaches? Has your head hurt?	Headache
Have you felt sick to your stomach like you were going to throw up?	Have you felt sick to your stomach or nauseous?	Nausea
Have you felt like you might fall when you walk, run or stand?	Have you had any balance problems or have you felt like you might fall	Balance problems
	when you walk, run or stand?	
Have you felt dizzy? (like things around you were spinning or moving)	Have you felt dizzy? (like things around you were spinning or moving)	Dizziness
Have you felt more tired than usual?	Have you felt more tired than usual?	Fatigue
Have bright lights bothered you more than usual? (like when you were in the sunlight,	Have bright lights bothered you more than usual? (like when you were in	Sensitivity to light
when you looked at lights, or watched TV)	the sunlight, when you looked at lights, or watched TV)	
Have loud noises bothered you more than usual? (like when people were talking,	Have loud noises bothered you more than usual (like when people were	Sensitivity to noise
when you heard sounds, watched TV or listened to loud music)	talking, when you heard sounds, watched TV or listened to loud music)	
Have you felt grumpy? (like you were in a bad mood)	Have you felt grumpy or irritable? (like you were in a bad mood)	Irritability
Have you felt sad?	Have you felt sad?	Sadness
Have you felt nervous or worried?	Have you felt nervous or worried?	Nervousness
Has it been hard for you to pay attention to what you are doing? (like homework or	Has it been hard for you to pay attention to what you are doing? (like	Difficulty concentrating
chores, listening to someone, or playing a game)	homework or chores, listening to someone, or playing a game)	
Has it been hard for you to remember things? (like things you heard or saw, or places	Has it been hard for you to remember things? (like things you heard or saw,	Difficulty remembering
you have gone)	or places you have gone)	
Have things looked fuzzy or blurry?	Have things looked blurry?	Vision problems
	Uncommon Elements Not Included	
Do you feel "different than usual?	Do you feel "different" than usual?	Feeling more emotional
	Have you felt like you are moving more slowly	Feeling slowed down
	Have you felt like you are thinking more slowly?	Feeling mentally foggy
	Has it been hard to think clearly?	Get confused with directions or tasks
	Have you felt more drowsy or sleepy than usual?	Move in a clumsy manner
		Answer questions more slowly than usual
		In general, to what degree do you feel
		"differently" than before the injury (not
		feeling yourself)

Supplemental Table 2: Comparison of patients included in heatmap vs. those missing data who were excluded from heatmap.

Characteristic	Included Patients	Excluded Patients	р
	N=2533	N=708	
Age, median (IQR)	12.0 (9.3-14.6)	12.1 (8.9-14.9)	0.77
Male Sex, n/total (%)	1416/2354 (60.2%)	441/708 (62.3%)	0.31
Time from Injury to ED Triage (hours), median (IQR)	3.0 (1.5-12.6)	2.7 (1.4-8.3)	0.04
Mechanism of Injury, n/total (%)			0.60
Sports/Recreation	1614/2353 (68.6%)	461/696 (66.2%)	
Fall	659 (28.0%)	212 (30.5%)	
Motor Vehicle Crash	45 (1.9%)	12 (!.7%)	
Assault	32 (1.4%)	11 (1.6%)	
Other	3 (0.1%)	0 (0%)	
Loss of Consciousness, n/total (%)	304/2084 (14.6%)	91/628 (14.5)	0.95
Post Injury Seizure, n/total (%)	41/2348 (1.7%)	16/693 (2.3%)	0.34
Glasgow Coma Score, median (IQR)	15 (15-15)	15 (15-15)	0.47
Past History, n/total (%)			
Prior Treatment for Headache	400/2353 (17.0%)	118/696 (17.0%)	0.98
Personal History of Migraine	310/2342 (13.2%)	82/696 (11.8%)	0.32
Learning Disability	179/2347 (7.6%)	64/692 (9.2%)	0.17
Attention Deficit Hyperactivity Disorder	195/2345 (8.3%)	73/691 (10.6%)	0.07
Anxiety	188/2349 (8.0%)	49/696 (7.0%)	0.41
Depression	62/2350 (2.6%)	25/697 (3.6%)	0.19
Sleep Disorder	47/2348 (2.0%)	15/692 (2.2%)	0.79
Other Psychiatric Disorders	24/2327 (1.0%)	8/689 (1.2%)	0.77
Family History of Migraine, n/total (%)	310/2342 (13.2%)	82/696 (11.8%)	0.32
BESS Number of Tandem Stance Errors, median (IQR)	3 (1-8)	3 (1-10)	0.03
BESS Number of Double Leg Stand Errors, median (IQR)	0 (0-1)	0 (0-1)	0.25
PCSI Scores at Index ED Visit, median (IQR)			
Physical	1.50 (0.88-2.38)	1.62 (0.88-2.62)	0.07
Fatigue	2.00 (0.67-3.67)	2.00 (1.00-3.67)	0.30
Emotional	0.50 (0.00-1.50)	0.75 (0.00-1.75)	0.15
Cognitive	0.60 (0.00-2.00)	0.80 (0.00-2.00)	0.12

Abbreviations: Balance Error Scoring System (BESS), Post-Concussion Symptom Inventory (PCSI)

## Supplemental Table 1: PCSI Symptom Inventory By Age with Common Question 13 Common Items Across All Age Groups Highlighted

Age 5-7	Age 8-12	Age 13-18
	Common Elements Included	
Have you had headaches? Has your head hurt?	Have you had headaches? Has your head hurt?	Headache
Have you felt sick to your stomach like you were going to throw up?	Have you felt sick to your stomach or nauseous?	Nausea
Have you felt like you might fall when you walk, run or stand?	Have you had any balance problems or have you felt like you might fall	Balance problems
	when you walk, run or stand?	
Have you felt dizzy? (like things around you were spinning or moving)	Have you felt dizzy? (like things around you were spinning or moving)	Dizziness
Have you felt more tired than usual?	Have you felt more tired than usual?	Fatigue
Have bright lights bothered you more than usual? (like when you were in the sunlight,	Have bright lights bothered you more than usual? (like when you were in	Sensitivity to light
when you looked at lights, or watched TV)	the sunlight, when you looked at lights, or watched TV)	
Have loud noises bothered you more than usual? (like when people were talking,	Have loud noises bothered you more than usual (like when people were	Sensitivity to noise
when you heard sounds, watched TV or listened to loud music)	talking, when you heard sounds, watched TV or listened to loud music)	
Have you felt grumpy? (like you were in a bad mood)	Have you felt grumpy or irritable? (like you were in a bad mood)	Irritability
Have you felt sad?	Have you felt sad?	Sadness
Have you felt nervous or worried?	Have you felt nervous or worried?	Nervousness
Has it been hard for you to pay attention to what you are doing? (like homework or	Has it been hard for you to pay attention to what you are doing? (like	Difficulty concentrating
chores, listening to someone, or playing a game)	homework or chores, listening to someone, or playing a game)	
Has it been hard for you to remember things? (like things you heard or saw, or places	Has it been hard for you to remember things? (like things you heard or saw,	Difficulty remembering
you have gone)	or places you have gone)	
Have things looked fuzzy or blurry?	Have things looked blurry?	Vision problems
	Uncommon Elements Not Included	
Do you feel "different than usual?	Do you feel "different" than usual?	Feeling more emotional
	Have you felt like you are moving more slowly	Feeling slowed down
	Have you felt like you are thinking more slowly?	Feeling mentally foggy
	Has it been hard to think clearly?	Get confused with directions or tasks
	Have you felt more drowsy or sleepy than usual?	Move in a clumsy manner
		Answer questions more slowly than usual
		In general, to what degree do you feel
		"differently" than before the injury (not
		feeling yourself)

Supplemental Table 2: Comparison of patients included in heatmap vs. those missing data who were excluded from heatmap.

Characteristic	Included Patients	Excluded Patients	р
	N=2533	N=708	
Age, median (IQR)	12.0 (9.3-14.6)	12.1 (8.9-14.9)	0.77
Male Sex, n/total (%)	1416/2354 (60.2%)	441/708 (62.3%)	0.31
Time from Injury to ED Triage (hours), median (IQR)	3.0 (1.5-12.6)	2.7 (1.4-8.3)	0.04
Mechanism of Injury, n/total (%)			0.60
Sports/Recreation	1614/2353 (68.6%)	461/696 (66.2%)	
Fall	659 (28.0%)	212 (30.5%)	
Motor Vehicle Crash	45 (1.9%)	12 (!.7%)	
Assault	32 (1.4%)	11 (1.6%)	
Other	3 (0.1%)	0 (0%)	
Loss of Consciousness, n/total (%)	304/2084 (14.6%)	91/628 (14.5)	0.95
Post Injury Seizure, n/total (%)	41/2348 (1.7%)	16/693 (2.3%)	0.34
Glasgow Coma Score, median (IQR)	15 (15-15)	15 (15-15)	0.47
Past History, n/total (%)			
Prior Treatment for Headache	400/2353 (17.0%)	118/696 (17.0%)	0.98
Personal History of Migraine	310/2342 (13.2%)	82/696 (11.8%)	0.32
Learning Disability	179/2347 (7.6%)	64/692 (9.2%)	0.17
Attention Deficit Hyperactivity Disorder	195/2345 (8.3%)	73/691 (10.6%)	0.07
Anxiety	188/2349 (8.0%)	49/696 (7.0%)	0.41
Depression	62/2350 (2.6%)	25/697 (3.6%)	0.19
Sleep Disorder	47/2348 (2.0%)	15/692 (2.2%)	0.79
Other Psychiatric Disorders	24/2327 (1.0%)	8/689 (1.2%)	0.77
Family History of Migraine, n/total (%)	310/2342 (13.2%)	82/696 (11.8%)	0.32
BESS Number of Tandem Stance Errors, median (IQR)	3 (1-8)	3 (1-10)	0.03
BESS Number of Double Leg Stand Errors, median (IQR)	0 (0-1)	0 (0-1)	0.25
PCSI Scores at Index ED Visit, median (IQR)			
Physical	1.50 (0.88-2.38)	1.62 (0.88-2.62)	0.07
Fatigue	2.00 (0.67-3.67)	2.00 (1.00-3.67)	0.30
Emotional	0.50 (0.00-1.50)	0.75 (0.00-1.75)	0.15
Cognitive	0.60 (0.00-2.00)	0.80 (0.00-2.00)	0.12

Abbreviations: Balance Error Scoring System (BESS), Post-Concussion Symptom Inventory (PCSI)